

For Tight Blood Sugar Control

^R ***Gliplite-M1/M2***

Glimepiride 1 mg. / 2 mg. + Metformin Hydrochloride 500 mg. Tablets



- › Controls both Fasting & Post Prandial blood sugar
- › Offers Superior reduction in FPG, PPG & HbA1c than Glipizide, Glyburide¹
- › Offers extra pancreatic effect on glucose metabolism by²
 - Improving peripheral glucose uptake
 - Decreasing exogenous glucose production

GLIP



Offers Safe & reliable diabetes control

^R *Metvilda* [®]

Vildagliptin 50 mg + Metformin 500 mg

|| Win over Diabetes Everyday ||



- » Improves dose proportionate HbA1c profile than Teneagliptin & Sitagliptin ¹
- » Significant reduction in fasting & post prandial blood glucose level
- » Better Circadian Glycaemic control than Sitagliptin ²
- » Better reduction of MAGE than Sitagliptin ²
- » Helps improving insulin sensitivity
- » Offers 24 hrs action

MET





Manage Post Prandial Hyperglycaemia

^R **VOGDDEEP** ^R

Voglibose 0.3 mg + Repaglinide 1 mg

A Trustworthy combination in Post Prandial Hyperglycemia



- ◆ Inhibits the over work of pancreatic Beta cell
- ◆ Increases of Reversion of IGT to Normal Glucose tolerance
- ◆ Offers significant reduction in 30 days ¹
 - Post Prandial blood glucose by **61.67** mg/dl
 - Fasting blood glucose by **42.13** mg/dl
- ◆ Effective & safe for the management of PPHG in T2DM patients ¹

VOG



A perfect approach in treating Type 2 Diabetes

^{Rx} **DGTURN** [®] **M**

Dapagliflozin 10 mg + Metformin 500 mg

Perfect Approach with Favourable Efficacy & Safety

Offers complementary mechanism of action, pharmacodynamics, pharmacokinetics, clinical efficacy and safety.



- Helps more patients achieve the desired goal by reducing HbA1c level ¹
- Increase medication adherence and patient satisfaction with improve glycemic control
- Low risk of hypoglycemia with added benefit of weight reduction and modest BP reduction
- Reduces the risk of HF in patients with Type 2 Diabetes ²

Also Available





Hypertension & Edema associated CHF

^{Rx} TORIFLOW

Torsemide 10 mg + Spironolactone 50mg

Ensure improvements in symptoms, protection & survival rates



- Prevent & protect against arrhythmia & HF ¹
- Well tolerated treatment of sodium & fluid retention
Resulting from moderate to severe CHF
- Effectively reduces pulmonary congestion,
peripheral edema, Jugular venus pressure ²
- Helps preventing strokes, HA & kidney problems ³
- Improvement in survival rates

TOR

Also Available





In Hypertension & Post MI Patients

℞ **Deepmet 25/50 XL**
Metoprolol 25 mg/50 mg (Extended Release)



- › Produce remarkable reduction in systolic & diastolic BP with one daily dosing ¹
- › Preferred Beta Blocker POST MI ²
30% reduction in mortality
48% reduction in mortality of diabetic survivor
- › Reduce the risk of death in HF patients with diabetes, MI or with history of Hypertension ³
- › No effect on blood glucose, HbA1c & insulin resistance by therapy, no side effect on diabetic hypertensive subject

DME





In the treatment of **Angina & Ischemic heart disease**

^R **NICODEEP**

Trimetazidine - 35mg Tablets

Add to every heart beat



- Increases survival in stable angina¹
- Protect against HF & Cardiomyopathy²
- Helps improving LV Volume & ejection fraction²
- Critical support in Ischemic crisis
- Significantly reduces mortality²

Prescribe confidently in

- **Stable Angia**
- **Myocardial Infraction**
- **Coronary Insufficiency**
- **IHD**

NIC






ACS, MI, Angina & Stroke

Rosedeep[®] GOLD 10/20

Rosuvastatin 10 mg / 20mg+ Clopidogrel 75 mg + Aspirin 75 mg Capsules

Synergistic Effect for Combined Benefits

Rosuvastatin: The Most Effective Statins in reducing LDL-C & Increasing HDL-C



Clopidogrel + Aspirin: Recommended by ACC, AHA & ACCP in ACS



- ♥ Acts in complimentary to each other with different mechanism
- ♥ Offers Synergism of stronger antiplatelet effect
- ♥ Superior antithrombotic protection in Coronary, Cerebral & Peripheral arterial disease

More patients reaches the Desired LDL-C reduction goal compared to Atorvastatin

	Rosuvastatin	Atorvastatin
NCEP ATP- III	80%	63%
Joint European Societies	82%	51%





In Hypertension

PROSARTAN

Telmisartan 40 mg


Tablets

A Superior ARB

Endorsed as First choice ARB
in Diabetes & Hypertension



Offers Superior 24 Hrs BP Controls¹



Offers Best M/E Ratio compared to
Losartan, Valsartan & Candesartan


In Stage II Hypertension
uncontrolled with Monotherapy

PROSARTAN-H

Telmisartan 40 mg + Hydrochlorothiazide 12.5 mg Tablets

A Trusted Combination

- ♥ Ensures 24 Hrs tight BP Control
- ♥ Offers Superior reduction of SBP & DBP than monotherapy alone²



Helps preventing strokes, heart attacks & kidney problems

Also Available



1. Journal of the American Society of Hypertension, Volume 2, Issue 4, Supplement, July-August 2008, Pages S16-S22
 2. American Journal of Therapeutics 9(2):111-7 March 2002 9(2):111-7






During Kidney dysfunction ensure recovery with

^R Creatlog™

Alpha Ketoanalogue

Support the Kidney to excrete urea



- Delays progression of CKD without deteriorating nutritional status¹
- Reduces the risk of commencing long term dialysis²
- Reduces uremic intoxication, proteinuria and amino aciduria, hyperphosphatemia
- Improves tubular functions, nutritional functions³

Prescribe confidently in

- Chronic Kidney Disease
- Kidney Dysfunction
- Diabetic Nephropathy

1 Tab BID

CRE





To relieve neuropathic pain, 45% of the patients requires two or more drugs for their pain relief



REJUDEEP[®]-NT

Pregabalin 75 mg + Methylcobalamin 1500 mcg + Nortriptyline 10 mg Tablets

NICE National Institute for Health and Care Excellence

Cited by NICE in the treatment strategies of Uncontrolled Neuropathic Pain

Highly recommended with highest grading by Danish expert committee



- Efficacious and safe therapy in the management of neuropathic pain
- Improves depression, mood, sleep disturbances, walking ability & overall health related quality of life

- Diabetic Neuropathy
- PIVD
- Low Back & Neck Pain
- Sciatica
- Neuralgia
- Radiculopathy
- Stroke & Fibromyalgia

1 Tab OD
At Night Time

Also Available
REJUDEEP-D
Pregabalin SR 75mg + Duloxetine 30mg





For the prevention **Thrombotic Events**

^R **TICODEEP-90**

Ticagrelor 90 mg



A Cornerstone therapy in ACS management



Recommended use of Ticagrelor for maintenance therapy in

- ▶ Patient with ACS (NSTEMI/STEMI) treated with DAPT after coronary stent implant
- ▶ Patient with NSTEMI treated with medical therapy alone

- ♥ Demonstrate faster onset & more potent inhibition of platelet aggregation than clopidogrel ¹
- ♥ Significantly reduces mortality rates from vascular causes, nonfatal MI & Strokes compared with clopidogrel ²
- ♥ Superior to clopidogrel in patient of ischemic event in patients with ACS ³
- ♥ Efficiency well tested & proved by PEGASUS TIMI, REDUCE-MVI, PLATO study

1. <https://www.drugs.com/medical-answers/ticagrelor-clopidogrel-354194>. 2. Heart. 2014 Nov;100(22):1762-9.
3. Catheter Cardiovasc Inter. 2022 May;99 Suppl 1:1395-1402





In Management of Hypertension with Diabetes or CKD


^{Rx} Tosles

Cilnidipine 10 mg Tablets


Better Control, Safer Control

Better Control Compared to Amlodipine

Significantly better daytime & nighttime BP control¹



Significantly greater reduction in pulse rate¹



- ♥ BP Control without reflex tachycardia²
- ♥ No significant pedal edema³
- ♥ Decrease proteinuria, improves renal function⁴

1. Hypertens Res 2005; 28: 1003-1008. 2. Hypertens Res 2005 Dec; 28(12):1003.8. 3. J Am Soc Hypertens. 2011 Sep-Oct; 5(5):410-6. 4. J Int Med Res. 2012;40(4):1417-28.



INDIA, the Diabetic Capital of World, in Every Diabetic patient


DHPL[®]-FOL

Methylcobalamin, L-Methylfolate & Pyridoxal - 5 - Phosphate Tablets

Reduces Homocysteine, Reduces Complications


Hyperhomocystenemia & Diabetes ¹

A dangerous association



**Exponentially increases
the risk of mortality upto **1000%****

**1 Tab OD for
4 to 6 Weeks**



Indian Journal of Clinical Biochemistry confirms:

Mecobalamin + Folic Acid + Pyridoxine



**Works synergistically and reduces
Hcy level in four to six weeks**

**Biologically Active, Highly Bio-available HLVs
Ensure Superior Results**

- Regenerate Nerves - Increases epidermal nerve fiber density
- Reduces Spontaneous Pain & Numbness of Limb by **73%** & **75%** ²
- Reduces the Risk of CVD & Stroke by **25%** ³

R

Diabetic Neuropathy

Hyperhomocysteinemia